

11/00 KAL

FILE 'MEDLINE, BIOSIS' ENTERED AT 15:51:16 ON 26 NOV 2000
L1 605566 S OSTEOCLAST# OR OSTEOBLAST# OR BONE
L2 65551 S ((INTERFERON OR INF) (A) (B OR G OR BETA OR GAMMA)) OR (INFB
OR
L3 2744 S L1 AND L2
L4 1241 S L3 AND (CANCER OR TUMOR OR MYELOMA OR METASTA? OR
CARCINOMA#)
L5 154 S L3 AND TUMOUR
L6 1280 S L4 OR L5
L7 808 S L6 AND HUMAN
L8 504 S L7 AND PY<1996
L9 217888 S L1/TI
L10 823 S L9 AND L2
L11 340 S L10 AND (CANCER OR TUMOR OR TUMOUR OR MYELOMA OR METASTA?
OR
L12 236 DUP REM L11 (104 DUPLICATES REMOVED)
L13 138 S L12 AND PY<1996
L14 14007 S OSTEOCLAST#
L15 265428 S METASTATIC OR METASTASIS
L16 538 S L15 AND L14
L17 3427 S L14(S) (INCREASE OR ACTIVATION OR UPREGULATION OR
(UP-REGULATI
L18 126 S L17 AND L15
L19 89 DUP REM L18 (37 DUPLICATES REMOVED)

National Library of Medicine: IGM Full Record Screen

**Related Articles**

TITLE: Nitric oxide and bone.

AUTHORS: Evans DM; Ralston SH

AUTHOR AFFILIATION: Department of Medicine and Therapeutics, University of Aberdeen, Scotland, UK.

SOURCE: J Bone Miner Res 1996 Mar;11(3):300-5

CITATION IDS: PMID: 8852940 UI: 97005641

ABSTRACT: Nitric oxide (NO), a mediator of cardiovascular homeostasis, neurotransmission, and immune function, has recently been found to have important effects in bone. Both constitutive and inducible forms of NO synthase are expressed by bone-derived cells, and cytokines such as interleukin-1 (IL-1), tumor necrosis factor (TNF), and interferon gamma (IFN-gamma), are potent stimulators of NO production. When combined with other cytokines, IFN-gamma markedly induces NO production, which suppresses osteoclast formation and activity of mature osteoclasts. This "superinduction" of NO is largely responsible for the selective inhibitory effect of IFN-gamma on cytokine-induced bone resorption. High concentrations of NO are also inhibitory for cells of the osteoblast lineage, and NO production appears to be partly responsible for the inhibitory effects of cytokines on osteoblast proliferation. At lower concentrations, however, NO has different effects. Moderate induction of NO potentiates bone resorption, and the constitutive production of NO at low concentrations promotes the proliferation of osteoblast-like cells and modulates osteoblast function. NO therefore appears to be an important regulatory molecule in bone with effects on cells of the osteoblast and osteoclast lineage and represents one of the molecules produced by osteoblasts which directly regulate osteoclastic activity. Stimulation of NO production in bone by proinflammatory cytokines raises the possibility that NO may be involved as a mediator of bone disease in conditions associated with cytokine activation, such as rheumatoid arthritis, tumor associated osteolysis, and postmenopausal

osteoporosis.

MAIN MESH HEADINGS: Bone and Bones/*metabolism
Nitric Oxide/*biosynthesis

ADDITIONAL MESH HEADINGS: Arthritis, Rheumatoid
Bone Resorption/chemically induced
Enzyme Induction/drug effects
Female
Free Radicals
Human
Interferon Type II/metabolism
Interferon Type II/pharmacology
Interleukin-1/metabolism
Interleukin-1/pharmacology
Nitric Oxide/metabolism
Nitric-Oxide Synthase/metabolism
Osteoblasts/cytology
Osteoblasts/metabolism
Osteoclasts/cytology
Osteoclasts/metabolism
Osteoporosis, Postmenopausal/etiology
Support, Non-U.S. Gov't
Tumor Necrosis Factor/metabolism
Tumor Necrosis Factor/pharmacology
1996/03
1996/01 00:00

PUBLICATION TYPES: JOURNAL ARTICLE
REVIEW
REVIEW, TUTORIAL

CAS REGISTRY NUMBERS: EC 1.14.13.39 (Nitric-Oxide Synthase)
0 (Free Radicals)
0 (Interleukin-1)
0 (Tumor Necrosis Factor)
10102-43-9 (Nitric Oxide)
82115-62-6 (Interferon Type II)

LANGUAGES: Eng



Order Documents



Log off IGM



Next Record



Details of Search



Return to Results



Return to Search Screen



Previous Record